

# Placebo Effects on the Neurologic Pain Signature

## A Meta-analysis of Individual Participant Functional Magnetic Resonance Imaging Data

Matthias Zunhammer, PhD; Ulrike Bingel, MD; Tor D. Wager, PhD; for the Placebo Imaging Consortium

**IMPORTANCE** Placebo effects reduce pain and contribute to clinical analgesia, but after decades of research, it remains unclear whether placebo treatments mainly affect nociceptive processes or other processes associated with pain evaluation.

**OBJECTIVE** We conducted a systematic, participant-level meta-analysis to test the effect of placebo treatments on pain-associated functional neuroimaging responses in the neurologic pain signature (NPS), a multivariate brain pattern tracking nociceptive pain.

**DATA SOURCES** Medline (PubMed) was searched from inception to May 2015; the search was augmented with results from previous meta-analyses and expert recommendations.

**STUDY SELECTION** Eligible studies were original investigations that were published in English in peer-reviewed journals and that involved functional neuroimaging of the human brain with evoked pain delivered under stimulus intensity-matched placebo and control conditions. The authors of all eligible studies were contacted and asked to provide single-participant data.

**DATA EXTRACTION AND SYNTHESIS** Data were collected between December 2015 and November 2017 following the Preferred Reporting Items for Systematic Review and Meta-Analyses of individual participant data guidelines. Results were summarized across participants and studies in a random-effects model.

**MAIN OUTCOMES AND MEASURES** The main, a priori outcome was NPS response; pain reports were assessed as a secondary outcome.

**RESULTS** We obtained data from 20 of 28 identified eligible studies, resulting in a total sample size of 603 healthy individuals. The NPS responses to painful stimulation compared with baseline conditions were positive in 575 participants (95.4%), with a very large effect size ( $g = 2.30$  [95% CI, 1.92 to 2.69]), confirming its sensitivity to nociceptive pain in this sample. Placebo treatments showed significant behavioral outcomes on pain ratings in 17 of 20 studies (85%) and in the combined sample ( $g = -0.66$  [95% CI,  $-0.80$  to  $-0.53$ ]). However, placebo effects on the NPS response were significant in only 3 of 20 studies (15%) and were very small in the combined sample ( $g = -0.08$  [95% CI,  $-0.15$  to  $-0.01$ ]). Similarly, analyses restricted to studies with low risk of bias ( $g = -0.07$  [95% CI,  $-0.15$  to  $0.00$ ]) indicated very small effects, and analyses of just placebo responders ( $g = -0.22$  [95% CI,  $-0.34$  to  $-0.11$ ]) indicated small effects, as well.

**CONCLUSIONS AND RELEVANCE** Placebo treatments have moderate analgesic effects on pain reports. The very small effects on NPS, a validated measure that tracks levels of nociceptive pain, indicate that placebo treatments affect pain via brain mechanisms largely independent of effects on bottom-up nociceptive processing.

JAMA Neurol. doi:10.1001/jamaneurol.2018.2017  
Published online July 30, 2018.

 Editorial

 Supplemental content

**Author Affiliations:** Klinik für Neurologie, Universitätsklinikum Essen, Essen, Germany (Zunhammer, Bingel); Department of Psychology and Neuroscience, University of Colorado, Boulder (Wager).

**Group Information:** Placebo Imaging Consortium members are listed at the end of this article.

**Corresponding Author:** Matthias Zunhammer, PhD, Klinik für Neurologie, Universitätsklinikum Essen, Hufelandstr. 55, 45147 Essen, Germany (matthias.zunhammer@uk-essen.de).

Placebo treatments are treatments with no intrinsic physical or pharmacological benefit. Nevertheless, they affect symptoms and physiology through patients' conceptions of the therapeutic context.<sup>1,2</sup> Placebo effects can be elicited by sham treatments, but they are also a substantial and beneficial part of the overall response to verum treatments, including those involving drugs and/or surgery.<sup>3,4</sup> In addition to conferring clinical benefits (and harms, in the case of nocebo effects), placebo effects reduce effect sizes for drug vs placebo differences in clinical trials, which may cause an increasing number of trials to fail<sup>5</sup> and thereby impede drug development.<sup>6,7</sup> Thus there is an urgent need to better understand placebo effects and to develop biomarkers for active drug responses, as well as placebo responses, to improve decision making in early clinical trials.<sup>8</sup>

Though placebo treatments can affect a variety of clinical outcomes,<sup>9</sup> placebo analgesia is the most robust and well studied.<sup>7,10,11</sup> Placebo analgesia has been linked with multiple psychological processes, including expectations and beliefs,<sup>12</sup> associative learning,<sup>13</sup> and social cognition.<sup>14</sup> Neurophysiological studies have suggested the involvement of descending inhibition of nociceptive afferents, with some studies supporting influences on spinal mechanisms<sup>15-19</sup> and others supporting higher-level cortical effects that cannot be explained by nociceptive input modulation alone, indicating affective or evaluative mechanisms.<sup>20-23</sup>

Nevertheless, the mechanisms by which placebo treatments change the perception of pain remain poorly understood, in part because of 2 limitations. First, previous studies were based on small sample sizes. Second, many brain areas associated with placebo analgesia, such as the anterior midcingulate cortex, the insula, and limbic regions, are involved in a range of functions, including cognitive decision making,<sup>24</sup> motor processes,<sup>25</sup> and emotion.<sup>26,27</sup> Thus, previous studies have not been able to establish whether placebo analgesia affects nociception-associated and pain-associated processing specifically or other cognitive and affective processes associated with the multidimensional experience of pain.

Recently, studies have begun to identify patterns of functional magnetic resonance imaging activity that yield objective and reliable<sup>28,29</sup> brain measures associated with evoked pain.<sup>30-33</sup> While they do not measure pain, which is by definition a subjective experience, they capture neurophysiological patterns associated with specific aspects of pain with high sensitivity.<sup>34,35</sup> Among ongoing efforts, the neurologic pain signature (NPS)<sup>33</sup> is a measure that has been shown to reliably track the intensity of evoked experimental pain across multiple studies with high sensitivity while responding only minimally to nonpainful somatic stimuli and other salient, aversive events, thus exhibiting high specificity.<sup>35-39</sup> Although fully understanding the neurophysiological processes captured by the NPS is a matter of ongoing investigation, previous results suggest that the NPS predominantly reflects changes in nociceptive input and the pain that arises from it, while being insensitive to higher cognitive pain modulation.<sup>36-39</sup>

In this study, we harness these methodological advances in a systematic meta-analysis of single-participant data testing placebo effects on NPS responses. If placebo treatments

## Key Points

**Question** How do placebo treatments affect pain processing in the brain?

**Findings** This systematic meta-analysis of single-participant functional magnetic resonance imaging data of 603 healthy participants from 20 studies found that placebo treatments against experimental pain have moderate effects on pain reports, but very small effects on the neurologic pain signature, a cerebral measure of nociceptive pain.

**Meaning** Placebo analgesia seems to be predominantly mediated by networks different from those underlying the primary processing of noxious stimuli.

predominantly affected early nociceptive processes, they would be expected to reduce activity in the NPS. If so, placebo effects and the endogenous pain-regulatory processes they engage may have pervasive effects on pain generation, making it hard to dissociate pain-associated and placebo-associated processes and outcomes. Conversely, if placebos mainly affect later-stage affective and evaluative processes, they may have little influence on NPS responses. In this case, it may be possible to develop meaningful measures of nociception that are placebo insensitive.

## Methods

We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses of Individual Participant Data.<sup>40</sup> The study protocol and hypotheses were registered (<https://osf.io/n9mb3/>) on December 10, 2015, which was after eligibility criteria and study search were defined, but before data collection and analysis began.

### Eligibility Criteria and Study Search

Criteria for study eligibility were peer-reviewed publication in the English language of an original investigation involving human participants who underwent functional neuroimaging of the brain during evoked pain and pain delivered under stimulus intensity-matched placebo and control conditions. Placebo treatment was defined as any condition where the experimental context suggested that an effective analgesic treatment was applied, including verbal suggestions and conditioning procedures that reinforced participants' expectations of reduced pain,<sup>41</sup> following the categorization of placebo paradigms introduced in Wager and Atlas in 2015.<sup>2</sup> Accordingly, nonplacebo control conditions that involved no treatment, ineffective treatment, hidden treatment (in contrast to open treatment), and unconditioned treatment (in contrast to conditioned treatment) were considered eligible. We identified potentially eligible studies on Medline on May 21, 2015 and December 6, 2015. This search was augmented by results from previous meta-analyses<sup>2,42</sup> and by requests for authors of individual placebo studies to identify additional studies missed in the online search. All 3 authors (M.Z., U.B., and T.D.W.) screened the titles and abstracts of

all records retrieved; studies that provisionally met eligibility criteria were assessed for eligibility by examining the full text. (Details appear in the eMethods and eTable 1 in the Supplement.)

### Data Acquisition

Authors of all eligible studies were contacted and asked to contribute participant-level neuroimaging, behavioral data (pain report), and demographic data. We requested data from the final, published analyses. Substudies with distinct participant samples and methods were treated as independent (studies 1 and 2 in Wager et al<sup>43</sup>). Reanalyses were excluded, but in the case of reanalyses with extended samples,<sup>44,45</sup> we included the study with the largest sample. Studies with partially missing data were included and analyzed based on the available data.<sup>46,47</sup>

### Outcome Definition

We obtained NPS responses<sup>33</sup> for each individual participant in each experimental condition and contrasted pain and placebo with pain and nonplacebo control conditions. The NPS responses were calculated as the dot product of each image with the NPS pattern, yielding a weighted average of activity across the image, where the NPS specifies the weights.<sup>33</sup> High NPS responses reflect both higher similarity of the individual images with the NPS pattern and higher-magnitude functional magnetic resonance imaging activity in the specified pattern.

To keep baseline conditions of studies with within-participant crossover designs and between-group placebo manipulations comparable, we limited our analysis to posttreatment conditions; in other words, additional baseline measurements from within-participant or mixed-design studies were excluded.<sup>46,48-50</sup> Further, nonpainful<sup>51</sup> or low-intensity<sup>43,46,48,52-54</sup> stimulus conditions were excluded from analysis to maximize detection sensitivity for placebo effects.

In several studies, images were provided for separate subconditions within placebo and control categories (eg, for left-lateralized and right-lateralized stimulation,<sup>55</sup> for strong and weak placebo conditions,<sup>56,57</sup> or for early-heat-pain and late-heat-pain periods).<sup>16,57,58</sup> In these cases, we summarized the NPS responses by calculating an average response under placebo and under control treatment for each participant (details appear in eTable 2 in the Supplement).

The scale of NPS responses depends on the scale of underlying imaging data and therefore on the image acquisition and analysis parameters used in the original studies. To avoid scaling issues, we based our analysis on the standardized effect size measure Hedges *g*, as is common in meta-analyses.<sup>59</sup> Similarly to the Cohen *d*, Hedges *g* is based on the mean difference between conditions divided by standard deviation, but with an additional correction for small sample bias. For within-participant studies we used Hedges  $g_{\text{im}}$ , which is based on the SD of within-participant differences corrected for within-participant correlations.<sup>60,61</sup>

### Risk of Bias Assessment

We used the Cochrane risk of bias tool<sup>62</sup> to evaluate the risk of bias for studies included in the present meta-analysis

(details are in the eMethods and eTable 3 in the Supplement). We assessed biases from selection (which arises via insufficient randomization), performance (via insufficient blinding of participants or treatment providers), detection (via insufficient blinding of analysts), attrition (by missing data), reporting (via underreporting of nonsignificant studies), and sequence (which is potentially introduced by within-participant designs).

### Analysis

Our main analysis followed a 3-part strategy. First, we tested the research question using all data available. Second, we conducted a conservative analysis excluding studies with high risk of bias. Third, we performed a responder analysis, in which we (1) only included participants who showed a behavioral placebo response (a pain report under placebo condition compared with the report under the control condition) above the study median, (2) excluded any experimental subconditions that may have diminished placebo effects (eg, placebo conditions deemed to be low efficacy or placebo conditions tested under pharmacological modulation), and (3) excluded any participants who were suspected outliers (the eMethods and eTable 2 in the Supplement).

Effects were summarized across studies using the generic inverse-variance weighting method with DerSimonian and Laird random effects,<sup>60</sup> meaning studies were weighted by  $1/SE^2$  (where SE is the standard error). We estimated heterogeneity in results using the  $\tau$  statistic, which represents the standard deviation of effect sizes between studies.<sup>59</sup> We tested against the null hypothesis of no effect at an error level of  $\alpha < .05$  (2-tailed), with additional inferences based on Bayes factors.<sup>63</sup> In brief, Bayes factors represent the relative likelihood of the null and the alternative hypotheses and have the advantage (compared with *P* values) that support for the null hypothesis can be concluded. Further analysis details and procedures used to check image quality are provided in the eMethods in the Supplement. Analysis was completed with MATLAB 2016b (MathWorks). The analysis code is available at <https://github.com/mzunhammer/PlaceboImagingMetaAnalysis>.

## Results

### Data Acquisition

We identified 96 published articles, of which 28 were selected as eligible. These included a total of 759 participants. Data from 20 studies and 603 participants were obtained, thus including the majority of eligible studies (71%) and participants (79.4%) published until 2015. Details on data acquisition are provided in eFigure 1 and eTable 1 in the Supplement. Eligible studies that could not be obtained were generally similar to the studies obtained in terms of pain stimulation, placebo induction, and study design; notably, 8 eligible studies with patient samples could not be obtained (eTable 4 in the Supplement). An update of the study search in March 2018 indicated that at least 6 eligible studies (with a combined patient population of 196 individuals) were published after data collection (in eMethods and eTable 1 in the Supplement). Even

Table. Included Studies

Source	Participants, No.	Design	Mean Age, y	Male, No. (%)	Pain Stimulus	Treatment Type	Placebo Suggestions	Placebo Conditioning	Placebo Comparison
Atlas et al, <sup>52</sup> 2012	21	Within	25	10 (48)	Contact heat	Remifentanyl infusion	Yes	No	Open vs hidden
Bingel et al, <sup>55</sup> 2006	19	Within	24	15 (79)	Laser	Inert topical cream	Yes	Yes	Effective vs ineffective
Bingel et al, <sup>49</sup> 2011	22	Within	28	15 (68)	Contact heat	Remifentanyl infusion	Yes	Yes	Open vs hidden
Choi et al, <sup>56</sup> 2011	15	Within	25	15 (100)	Electrical	Saline infusion	Yes	Yes	Effective vs ineffective
Eippert et al, <sup>16</sup> 2009	40	Within	26	40 (100)	Contact heat	Inert topical cream	Yes	Yes	Effective vs ineffective
Ellingsen et al, <sup>51</sup> 2013	28	Within	26	19 (68)	Contact heat	Inert nasal spray	Yes	No	Treatment vs no treatment
Elsenbruch et al, <sup>64</sup> 2012	36	Within	26	15 (42)	Distension	Saline infusion	Yes	No	Effective vs ineffective
Freeman et al, <sup>53</sup> 2015	24	Within	27	12 (50)	Contact heat	Inert topical cream	Yes	Yes	Effective vs ineffective
Geuter et al, <sup>57</sup> 2013	40	Within	26	40 (100)	Contact heat	Inert topical cream	Yes	Yes	Effective vs ineffective
Kessner et al, <sup>50</sup> 2013	39	Between	26	20 (51)	Contact heat	Inert topical cream	No	Yes	Effective vs ineffective
Kong et al, <sup>46</sup> 2006	10 <sup>c</sup>	Within	27	6 (60)	Contact heat	Sham acupuncture	Yes	Yes	Effective vs ineffective
Kong et al, <sup>48</sup> 2009	12 <sup>a</sup>	Within	26	5 (42)	Contact heat	Sham acupuncture	Yes	Yes	Effective vs ineffective
Lui et al, <sup>65</sup> 2010	31	Within	23	15 (45)	Laser	Sham tens	Yes	Yes	Effective vs ineffective
Rütgen et al, <sup>54</sup> 2015	102	Between	25	32 (31)	Electrical	Inert pill	Yes	Yes	Treatment vs no treatment
Schenk et al, <sup>79</sup> 2014	32	Within	26	17 (53)	Capsaicin and heat	Topical lidocaine and inert topical cream	Yes	No	Hidden vs open (lidocaine), effective vs ineffective (inert cream)
Theysohn et al, <sup>45</sup> 2014	30	Within	35	15 (50)	Distension	Saline infusion	Yes	No	Effective vs ineffective
Wager et al, <sup>43</sup> 2004a <sup>b</sup>	24	Within	na	na	Electrical	Inert topical cream	Yes	No	Effective vs ineffective
Wager et al, <sup>43</sup> 2004b <sup>c</sup>	23	Within	na	na	Contact heat	Inert topical cream	Yes	Yes	Effective vs ineffective
Wrobel et al, <sup>58</sup> 2014	38	Within	26	22 (58)	Contact heat	Inert topical cream	Yes	Yes	Effective vs ineffective
Zeidan et al, <sup>47</sup> 2015	17 <sup>a</sup>	Within	28	8 (47)	Contact heat	Inert topical cream	Yes	Yes	Treatment vs no treatment

Abbreviations: NA, not available; TENS, transcutaneous electrical nerve stimulation.

<sup>a</sup> Placebo-treatment groups only.

<sup>b</sup> Substudy 1.

<sup>c</sup> Substudy 2.

when considering these additional studies, the present meta-analysis covers most eligible studies (59%) and participants (63.1%).

### Risk of Bias

For pain ratings, the assessment of risk of bias (eResults, eTable 3, and eFigure 2 in the [Supplement](#)) indicated a high risk of performance (self-report) and detection bias, as well as unknown levels of reporting (publication) bias. For NPS responses, we found low risk of bias, because this measure does not depend on self-report and was unknown when the original studies were performed.

### Sample Description


Included studies are listed in the [Table](#), and key sample characteristics are shown in eFigure 3 in the [Supplement](#). Details on pain stimulation, placebo treatment, image

acquisition, and imaging analyses are provided in the eMethods and eTables 5, 6, 7, and 8 in the [Supplement](#). Image alignment to Montreal Neurological Institute space was satisfactory, and the coverage of the voxels making up the NPS was near optimum levels (98.4% across all participants; eMethods and eFigure 4 in the [Supplement](#)). Four participants showed mean pain ratings less than 5% of the pain scale, indicating insufficient pain stimulation; evidence for imaging artifacts was found in 12 participants (2.0%) (eMethods in the [Supplement](#)). These participants were defined as outliers and excluded from the responder analysis, but retained in the primary and the conservative analysis.

### NPS Responses to Painful Stimulation

The NPS responses to painful stimulation, compared with low-level baseline of placebo and control conditions pooled,

Figure 1. Neurologic Pain Signature Response to Noxious Stimulation vs Baseline in All Studies



were very large ( $g = 2.30$  [95% CI, 1.92 to 2.69]) and significant in all included studies (Figure 1). Positive NPS responses were observed in 576 individuals (95.4%) (95.6% when outliers were excluded). The estimated between-study standard deviation of effect sizes  $\tau$  was 0.68, indicating that the outcomes of painful stimulation on NPS responses varied considerably among studies, which was as expected, given the broad range of different imaging and pain stimulation protocols used (Table; eFigure 3 and eTables 5, 6, 7, and 8 in the Supplement). Moreover, the NPS responded more strongly to high vs low levels of stimulus intensity in 6 of 7 studies (86%) in which this comparison was possible ( $g = 1.32$  [95% CI, 0.86 to 1.79]; eFigure 5 in the Supplement). Together, these findings validate that the NPS was sensitive to noxious stimulation in this sample.

### Placebo Effects on Pain Ratings and NPS Responses

Placebo treatments, compared with matched control conditions, showed moderate<sup>66</sup> analgesic effects on pain ratings ( $g = -0.66$  [95% CI, -0.80 to -0.53]; Figure 2), which corresponds to a reduction of -11.3 (95% CI, -14.0 to -8.56) units on a 101-point visual analogue scale. The Bayes factor ( $B_{N(0, 0.5)}$ ) obtained for a normal null-prior ( $g = 0$  [SD, 0.5], 2-tailed) was  $9.4 \times 10^{18}$ , indicating overwhelming support for the hypothesis of nonnull placebo effects on pain ratings.<sup>63</sup> Effect sizes varied considerably among studies ( $\tau = 0.24$ ), which may be explained by the variation in placebo paradigms used (Table; eTable 6 in the Supplement).

In contrast, effects of placebo treatments on NPS responses were small ( $g = -0.08$  [95% CI, -0.15 to -0.01]; Figure 3), with little between-study heterogeneity ( $\tau = 0.015$ ). Thus, placebo effects on the NPS were 12.1% as

large as effects on pain ratings, and 3.7% as large as the effects of painful stimulation on the NPS. The Bayes factor was less than 1 ( $B_{N(0, 0.5)} = 0.805$ ), indicating that these data provide very weak support in favor of the null hypothesis of no effect.<sup>63</sup>


A conservative analysis excluding all studies with high risk of bias and therefore including 15 studies with 429 participants yielded similar results (placebo - control on NPS,  $g = -0.07$  [95% CI, -0.15 to 0.00];  $B_{N(0, 0.5)} = 0.787$ ; eFigure 6 in the Supplement). Somewhat larger effects were found in the responder analysis, which included only participants showing a behavioral placebo response greater than the study median and excluded potentially ineffective placebo treatments and outliers (196 participants from 18 studies were included). In this sample of so-called placebo responders, the Bayes factor ( $B_{N(0, 0.5)} = 113.8$ ) indicated robust support for the hypothesis of a nonnull placebo effect on NPS responses. However, the effects remained in the small range ( $g = -0.22$  [95% CI, -0.34 to -0.11]; eFigure 7 in the Supplement). Thus, even in placebo responders, effects of placebo on the NPS were only 4% to 14% as large as the overall NPS response to painful stimulation (Figure 1).

### Associations Between Placebo Effects on Pain Reports and NPS Responses

For studies with crossover designs, which included within-participant testing of both placebo and control treatments, we performed a meta-analysis of within-study correlations between placebo effects on pain report and NPS responses across individuals. A Bayes factor of 894.5 ( $B_{N(0, 0.5)}$ ) indicated robust support for the hypothesis of a nonnull




Figure 2. Changes in Pain Ratings After Experimental Placebo Treatment



Studies were weighted according to inverse variance. Total  $z = 9.65$  ( $P < .001$ ); heterogeneity  $\chi^2_{19} = 55.35$  ( $P < .001$ );  $\tau^2 = 0.06$ ;  $I^2 = 65.68\%$ .

Figure 3. Changes in Neurologic Pain Signature Responses After Experimental Placebo Treatments



Studies were weighted according to inverse variance. Total  $z = -2.19$  ( $P = .03$ ); heterogeneity  $\chi^2_{19} = 19.15$  ( $P = .45$ );  $\tau^2 = 0.00$ ;  $I^2 = 0.77\%$ .

correlation, with greater placebo analgesia associated with greater NPS downregulation (Pearson  $r = 0.23$  [95% CI, 0.13 to 0.33];  $P < .001$ ; eFigure 8 in the Supplement). While statistically significant, the effect was in the small to moderate range, suggesting that placebo analgesia is weakly associated with NPS downregulation.

### Comparing Placebo Effects on the NPS With Effect of Reduced Stimulus Intensity

These findings suggest that the effects of placebo are small in terms of effective changes in nociceptive input. To further quantify placebo effects on the NPS in terms of equivalent changes in noxious stimulus intensity, we compared placebo

effects with effects of fixed increases in physical stimulus intensity. For this purpose, we used a convenience sample of 3 available out-of-sample studies<sup>67-69</sup> that tested the effects of fixed increments in noxious heat (steps of 0.4°C in a range from 45.9°C to 47.5°C<sup>67</sup> and steps of 1°C in a range from 46°C to 48°C<sup>69</sup>) or noxious pressure (reduction of 1.5 kg/cm<sup>2</sup> from a baseline of 6.0 kg/cm<sup>2</sup>).<sup>68</sup> These independent studies were used because stimulus intensities in our meta-analysis sample varied across individuals and produced behavioral effects several times larger than those of placebo, making estimation of equivalent stimulus intensity differences unreliable (for example, compare Figure 2 and eFigure 5 in the [Supplement](#)). Effect sizes (*g*) for stimulus intensity on pain ratings were -0.63 (95% CI, -0.68 to -0.58) at -0.4°C, -1.10 (95% CI, -1.30 to -0.90) at -1.0°C, and -1.24 (95% CI, -1.66 to -0.81) at -1.5 kg/cm<sup>2</sup> (eFigure 9 in the [Supplement](#)), compared with -0.66 (95% CI, -0.80 to -0.53) for placebo.

Placebo effects on NPS responses in the full meta-analysis sample were 5.0 and 8.9 times smaller than the effect of reducing heat by -0.4°C<sup>67</sup> and -1.0°C,<sup>69</sup> respectively, and 10.9 times smaller than the effect of reducing painful nail bed pressure by -1.5 kg/cm<sup>2</sup>.<sup>68</sup> (eFigure 9 in the [Supplement](#)), corresponding to an effective reduction in noxious heat of approximately 0.1°C and in noxious pressure of approximately 0.14 kg/cm<sup>2</sup>. Thus, placebo effects on the NPS are small compared with modest reductions in stimulus intensity, despite similar effects on pain ratings.

### Comparing Placebo and Remifentanyl Effects on the NPS

Two of the studies in the sample used analgesic, nonsedative doses of the  $\mu$ -opioid agonist remifentanyl<sup>49,52</sup> at similar doses (with brain remifentanyl concentrations of 0.76 ng/mL<sup>52</sup> and 0.80 ng/mL<sup>49</sup>). This allowed comparison of the effects of opioids and placebo treatment on NPS responses within these studies. Remifentanyl (*g* = -0.77 [95% CI, -1.39 to -0.18]) and placebo treatments (*g* = -0.79 [95% CI, -1.39 to -0.18]) induced comparable analgesic effects at the behavioral level. However, the effects of remifentanyl on the NPS (-1.10 [95% CI, -1.44 to -0.76] *g*) were about 10 times larger than the mean effect of placebo treatments in these studies (*g* = -0.11 [95% CI, -0.38 to 0.16]; eFigure 10 in the [Supplement](#)). These estimates indicate that the effects of placebo treatment on NPS responses are small compared with the analgesic effects of opioids, despite comparable behavioral effects.

## Discussion

This large-scale meta-analysis of participant-level data revealed that placebo treatments have moderate effects on subjective reports of pain, but minimal effects on responses in the NPS, a central nervous system marker that tracks the intensity of nociceptive pain. These findings are based on most of the neuroimaging data on placebo analgesia published in the field until 2015. Results were consistent across a variety of pain-induction and placebo-induction methods and across 3 parallel analyses varying in risk of bias, including an analysis limited to placebo responders.

These results extend our understanding of placebo analgesia by suggesting that the effects of placebos on cerebral pain stimulus intensity processing are limited. We did observe small reductions in NPS response, which scaled with individual analgesia. Such effects may reflect descending inhibition of nociceptive systems, consistent with that findings of placebo, nocebo, and cognitive effects on spinal functional magnetic resonance imaging signals and brainstem nuclei involved in descending modulation that have been reported in previous studies.<sup>15,17-19,70</sup> However, the very small size of the effects on the NPS argues against a strong and pervasive early influence and point to stronger influences of other systems independent of the NPS. Our results emphasize that placebo analgesia is a phenomenon not based on a single mechanism but rather multiple mechanisms that have yet to be fully understood.

Importantly, the NPS is not a complete model of pain and pain-associated functionality and was not intended as such.<sup>33</sup> Lack of effects on the NPS does not imply that placebo effects do not influence pain perception or pain-associated behavior. Indeed, the insensitivity of the NPS to manipulations that affect reported pain<sup>35-39</sup> implies that there must be other processes that contribute to pain report. However, the high sensitivity of the NPS to variation in nociceptive input (details may be found in Figure 1; eFigure 5 and eFigure 9 in the [Supplement](#); and previous publications<sup>33,35-39</sup>), and its sensitivity to known analgesics (eg, the  $\mu$ -opioid agonist remifentanyl; eFigure 10 in the [Supplement](#)), suggest that if placebo treatments had pervasive early effects on pain processing, they should have been reflected in the NPS. They did not, leading us to infer that the placebo treatments studied here affect processes that are largely consequent to activation of nociceptive systems. Such processes include cognitive evaluation,<sup>71</sup> pain affect, pain-associated decisionmaking, and mesolimbic reward processing.<sup>72</sup> These processes are likely important for behavior and subjective well-being in their own right, and indeed, placebo treatments can impact long-term symptom perception and functionality in clinically meaningful ways,<sup>73,74</sup> whether they impact nociceptive pain signaling or not.

These findings also have implications for the objective assessment of treatment effects on pain-associated neurophysiology. Patients evaluate their pain within a complex set of personal and cultural factors,<sup>75</sup> which poses challenges for clinical trials that use self-reported pain as a primary outcome. Whether patients feel better is paramount for overall well-being, but it does not guarantee that a treatment impacts the intended physiological mechanisms in the brain and elsewhere in the body. Objective neurophysiological measures do not replace reported pain and well-being, but they can provide measures of pharmacodynamic efficacy on specific brain targets.<sup>8,76</sup> The present study further establishes the NPS as a brain measure that is sensitive to multiple types of evoked pain and insensitive to cognitive factors<sup>33,35-39</sup> in a large and geographically diverse sample. Because the NPS was found sensitive to opioid drugs but not placebo treatment, this may make it an appealing target for evaluating pharmacodynamics and efficacy in early-stage clinical trials.<sup>8,76</sup> Our findings suggest that brain patterns such as the NPS can be used to assess efficacy in modulating nociceptive systems for drugs or devices that are

intended to modulate nociceptive input at the peripheral or spinal level or influence descending facilitation or inhibition. As suggested by Duff et al,<sup>8</sup> the NPS and associated markers could be used to make early stop or go decisions in phase II clinical trials, before much more costly large-scale tests in phase III studies.

### Limitations

Several caveats deserve mention. First, this meta-analysis included only studies testing experimental placebo treatments for evoked pain in healthy participants. They may not generalize to clinical pain, which likely involves a complex mix of nociceptive and extranociceptive processes.<sup>77</sup> Second, we analyzed summary images from published analyses. While this helps to ensure careful quality control and is advantageous in ensuring broad generalizability of results, it likely increases interstudy heterogeneity and reduces overall effect sizes. However, these issues are unlikely to compromise our conclusions regarding placebo effects, because we compare them with strong positive controls. Third, the present meta-analysis only covers the relevant literature until mid-2015; more recent studies were not sought because of the time demands of collecting and analyzing participant-level imaging data. Advances in data sharing and standardization will hopefully make it possible to perform participant-level meta-analyses more quickly in the future.

Importantly, not all placebo manipulations are likely to be equally effective, as indicated by the heterogeneity in placebo effects on pain ratings (as shown in Figure 2 and described in previous publications<sup>10,11</sup>). Although the studies tested here were fairly homogenous in terms of placebo effect on the NPS (Figure 3), there are multiple pathways to cognitive pain modulation,<sup>37,78</sup> and some may affect the NPS more strongly than others. Treatment contexts not studied here may still influence NPS responses.

### Conclusions

In sum, we have shown that placebo treatments have only small effects on a cerebral pattern tracking nociceptive pain in what is to our knowledge the largest meta-analysis of single-participant neuroimaging data on this topic to date. This suggests that placebo analgesia is largely mediated by networks different from those underlying the primary processing of noxious stimuli. Further studies are necessary to better understand which aspects of pain processing are affected by placebo treatments, and the significance of those processes for long-term clinical outcomes and wellbeing. This work serves as a starting point for the development of brain models that track pain-associated outcomes and other clinical and behavioral endpoints.

### ARTICLE INFORMATION

**Accepted for Publication:** May 1, 2018.

**Author Contributions:** Drs Bingel and Wager had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** All authors.

**Acquisition, analysis, or interpretation of data:** All authors.

**Drafting of the manuscript:** All authors.

**Critical revision of the manuscript for important intellectual content:** All authors.

**Statistical analysis:** Zunhammer, Wager.

**Obtained funding:** Bingel, Wager.

**Administrative, technical, or material support:** Bingel.

**Supervision:** Bingel, Wager.

**Published Online:** July 30, 2018.

doi:10.1001/jamaneuro.2018.2017

**Group Information:** Consortium authors: Lauren Atlas, PhD, National Center for Complementary and Integrative Health, National Institutes of Health, Bethesda, Maryland, and National Institute on Drug Abuse, National Institutes of Health, Baltimore, Maryland; Fabrizio Benedetti, MD, University of Turin, Turin, Italy and Plateau Rosà Labs, Plateau Rosà, Switzerland; Christian Büchel, MD, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; Jae Chan Choi, MD, PhD, Yonsei University, Wonju, South Korea; Luana Colloca, MD, PhD, University of Maryland, Baltimore, Maryland; Davide Duzzi, PhD, University of Modena e Reggio Emilia, Modena, Italy; Falk Eippert, PhD, University of Oxford, Oxford, United Kingdom; Dan-Mikael Ellingsen, PhD, Harvard Medical School, Cambridge, Massachusetts; Sigrid Elsenbruch, PhD, Essen University Hospital, Essen, Germany; Stephan

Geuter, PhD, Johns Hopkins University, Baltimore, Maryland; Randy Gollub, MD, PhD, Harvard Medical School, Cambridge, Massachusetts;

Ted J. Kaptchuk, Beth Israel Deaconess Medical, Harvard Medical School, Boston, Massachusetts; Simon S. Kessner, MD, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; Irving Kirsch, PhD, Harvard Medical School, Cambridge, Massachusetts; Jian Kong, MD, Massachusetts General Hospital, Harvard Medical School, Cambridge, Massachusetts; Claus Lamm, PhD, University of Vienna, Vienna, Austria; Siri Leknes, PhD, University of Oslo, Oslo, Norway; Alexa Müllner-Huber, PhD, PhD, University of Vienna, Vienna, Austria; Fausta Lui, MD, University of Modena e Reggio Emilia, Modena, Italy; Carlo A. Porro, MD, PhD, University of Modena e Reggio Emilia, Modena, Italy; Markus Rütgen, PhD, University of Vienna, Vienna, Austria; Lieven Schenk, PhD, University of Maryland, Baltimore, Maryland; Julia Schmid, PhD, Essen University Hospital, Essen, Germany; Nina Theysohn, MD, Essen University Hospital, Essen, Germany; Irene Tracey, PhD, University of Oxford, Oxford, United Kingdom; Nathalie Wrobel, PhD, Karolinska Institute, Solna, Sweden; Fadel Zeidan, PhD, Wake Forest School of Medicine, Winston-Salem, North Carolina.

**Conflict of Interest Disclosures:** None reported.

**Funding/Support:** This research was funded by the German Research Foundation (FOR 1328), the Mercator Research Center Ruhr (MERCUR), and National Institutes of Health (grants R01 MH076136 and R01 DA035484, Dr Wager).

**Role of the Funder/Sponsor:** Funders had no influence on design and conduct of the study; collection, management, analysis, and

interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

### REFERENCES

1. Ashar YK, Chang LJ, Wager TD. Brain mechanisms of the placebo effect: an affective appraisal account. *Annu Rev Clin Psychol*. 2017;13:73-98. doi:10.1146/annurev-clinpsy-021815-093015
2. Wager TD, Atlas LY. The neuroscience of placebo effects: connecting context, learning and health. *Nat Rev Neurosci*. 2015;16(7):403-418. doi:10.1038/nrn3976
3. Khan A, Kolts RL, Rapaport MH, Krishnan KRR, Brodhead AE, Browns WA. Magnitude of placebo response and drug-placebo differences across psychiatric disorders. *Psychol Med*. 2005;35(5):743-749. doi:10.1017/S0033291704003873
4. Jonas WB, Crawford C, Colloca L, et al. To what extent are surgery and invasive procedures effective beyond a placebo response? A systematic review with meta-analysis of randomised, sham controlled trials. *BMJ Open*. 2015;5(12):e009655. doi:10.1136/bmjopen-2015-009655
5. Tuttle AH, Tohyama S, Ramsay T, et al. Increasing placebo responses over time in U.S. clinical trials of neuropathic pain. *Pain*. 2015;156(12):2616-2626. doi:10.1097/j.pain.0000000000000333
6. Borsook D, Hargreaves R, Bountra C, Porreca F. Lost but making progress—where will new analgesic drugs come from? *Sci Transl Med*. 2014;6(249):249sr3. doi:10.1126/scitranslmed.3008320
7. Enck P, Bingel U, Schedlowski M, Rief W. The placebo response in medicine: minimize, maximize or personalize? *Nat Rev Drug Discov*. 2013;12(3):191-204. doi:10.1038/nrd3923



8. Duff EP, Vennart W, Wise RG, et al. Learning to identify CNS drug action and efficacy using multistudy fMRI data. *Sci Transl Med*. 2015;7(274):274ra16. doi:10.1126/scitranslmed.3008438
9. Finniss DG, Kaptchuk TJ, Miller F, Benedetti F. Biological, clinical, and ethical advances of placebo effects. *Lancet*. 2010;375(9715):686-695. doi:10.1016/S0140-6736(09)61706-2
10. Hróbjartsson A, Gøtzsche PC. Placebo interventions for all clinical conditions (review) placebo interventions for all clinical conditions. *Cochrane Database Syst Rev*. 2010;(1):CD003974. doi:10.1002/14651858
11. Vase L, Petersen GL, Riley JL III, Price DD. Factors contributing to large analgesic effects in placebo mechanism studies conducted between 2002 and 2007. *Pain*. 2009;145(1-2):36-44. doi:10.1016/j.pain.2009.04.008
12. Atlas LY, Wager TD. How expectations shape pain. *Neurosci Lett*. 2012;520(2):140-148. doi:10.1016/j.neulet.2012.03.039
13. Zunhammer M, Ploner M, Engelbrecht C, Bock J, Kessner SS, Bingel U. The effects of treatment failure generalize across different routes of drug administration. *Sci Transl Med*. 2017;9(393):1-8. doi:10.1126/scitranslmed.aal2999
14. Colloca L, Benedetti F. Placebo analgesia induced by social observational learning. *Pain*. 2009;144(1-2):28-34. doi:10.1016/j.pain.2009.01.033
15. Goffaux P, Redmond WJ, Rainville P, Marchand S. Descending analgesia—when the spine echoes what the brain expects. *Pain*. 2007;130(1-2):137-143. doi:10.1016/j.pain.2006.11.011
16. Eippert F, Bingel U, Schoell ED, et al. Activation of the opioidergic descending pain control system underlies placebo analgesia. *Neuron*. 2009;63(4):533-543. doi:10.1016/j.neuron.2009.07.014
17. Matre D, Casey KL, Knardahl S. Placebo-induced changes in spinal cord pain processing. *J Neurosci*. 2006;26(2):559-563. doi:10.1523/JNEUROSCI.4218-05.2006
18. Geuter S, Büchel C. Facilitation of pain in the human spinal cord by nocebo treatment. *J Neurosci*. 2013;33(34):13784-13790. doi:10.1523/JNEUROSCI.2191-13.2013
19. Tinnermann A, Geuter S, Sprenger C, Finsterbusch J, Büchel C. Interactions between brain and spinal cord mediate value effects in nocebo hyperalgesia. *Science*. 2017;358(6359):105-108. doi:10.1126/science.aan1221
20. Wager TD, Matre D, Casey KL. Placebo effects in laser-evoked pain potentials. *Brain Behav Immun*. 2006;20(3):219-230. doi:10.1016/j.bbi.2006.01.007
21. Martini M, Lee MCH, Valentini E, Iannetti GD. Intracortical modulation, and not spinal inhibition, mediates placebo analgesia. *Eur J Neurosci*. 2015;41(4):498-504. doi:10.1111/ejn.12807
22. Petrovic P, Kalso E, Petersson KM, Andersson J, Fransson P, Ingvar M. A prefrontal non-opioid mechanism in placebo analgesia. *Pain*. 2010;150(1):59-65. doi:10.1016/j.pain.2010.03.011
23. Wager TD, Atlas LY, Leotti LA, Rilling JK. Predicting individual differences in placebo analgesia: contributions of brain activity during anticipation and pain experience. *J Neurosci*. 2011;31(2):439-452. doi:10.1523/JNEUROSCI.3420-10.2011
24. Grinband J, Hirsch J, Ferrera VP. A neural representation of categorization uncertainty in the human brain. *Neuron*. 2006;49(5):757-763. doi:10.1016/j.neuron.2006.01.032
25. Dum RP, Levinthal DJ, Strick PL. Motor, cognitive, and affective areas of the cerebral cortex influence the adrenal medulla. *Proc Natl Acad Sci U S A*. 2016;113(35):9922-9927. doi:10.1073/pnas.1605044113
26. Lindquist KA, Wager TD, Kober H, Bliss-Moreau E, Barrett LF. The brain basis of emotion: a meta-analytic review. *Behav Brain Sci*. 2012;35(3):121-143. doi:10.1017/S0140525X11000446
27. Wager TD, Atlas LY, Botvinick MM, et al. Pain in the ACC? *Proc Natl Acad Sci U S A*. 2016;113(18):E2474-E2475. doi:10.1073/pnas.1600282113
28. Woo C-W, Wager TD. Neuroimaging-based biomarker discovery and validation. *Pain*. 2015;156(8):1379-1381. doi:10.1097/j.pain.0000000000000223
29. Woo C-W, Wager TD. What reliability can and cannot tell us about pain report and pain neuroimaging. *Pain*. 2016;157(3):511-513. doi:10.1097/j.pain.0000000000000442
30. Marquand A, Howard M, Brammer M, Chu C, Coen S, Mourão-Miranda J. Quantitative prediction of subjective pain intensity from whole-brain fMRI data using Gaussian processes. *Neuroimage*. 2010;49(3):2178-2189. doi:10.1016/j.neuroimage.2009.10.072
31. Brodersen KH, Wiech K, Lomakina EI, et al. Decoding the perception of pain from fMRI using multivariate pattern analysis. *Neuroimage*. 2012;63(3):1162-1170. doi:10.1016/j.neuroimage.2012.08.035
32. Brown JE, Chatterjee N, Younger J, Mackey S. Towards a physiology-based measure of pain: patterns of human brain activity distinguish painful from non-painful thermal stimulation. *PLoS One*. 2011;6(9):e24124. doi:10.1371/journal.pone.0024124
33. Wager TD, Atlas LY, Lindquist MA, Roy M, Woo CW, Kross E. An fMRI-based neurologic signature of physical pain. *N Engl J Med*. 2013;368(15):1388-1397. doi:10.1056/NEJMoal204471
34. Reddan MC, Lindquist MA, Wager TD. Effect size estimation in neuroimaging. *JAMA Psychiatry*. 2017;74(3):207-208. doi:10.1001/jamapsychiatry.2016.3356
35. Woo C-W, Chang LJ, Lindquist MA, Wager TD. Building better biomarkers: brain models in translational neuroimaging. *Nat Neurosci*. 2017;20(3):365-377. doi:10.1038/nn.4478
36. Woo C-W, Roy M, Buhle JT, Wager TD. Distinct brain systems mediate the effects of nociceptive input and self-regulation on pain. *PLoS Biol*. 2015;13(1):e1002036. doi:10.1371/journal.pbio.1002036
37. Woo C-W, Schmidt L, Krishnan A, et al. Quantifying cerebral contributions to pain beyond nociception. *Nat Commun*. 2017;8:14211. doi:10.1038/ncomms14211
38. Becker S, Gandhi W, Pomares F, Wager TD, Schweinhardt P. Orbitofrontal cortex mediates pain inhibition by monetary reward. *Soc Cogn Affect Neurosci*. 2016;12(4):651-661. doi:10.1093/scan/nsw173
39. Bräscher A-K, Becker S, Hoeppli M-E, Schweinhardt P. Different brain circuitries mediating controllable and uncontrollable pain. *J Neurosci*. 2016;36(18):5013-5025. doi:10.1523/JNEUROSCI.1954-15.2016
40. Stewart LA, Clarke M, Rovers M, et al; PRISMA-IPD Development Group. Preferred reporting items for systematic review and meta-analyses of individual participant data: the PRISMA-IPD statement. *JAMA*. 2015;313(16):1657-1665. doi:10.1001/jama.2015.3656
41. Voudouris NJ, Peck CL, Coleman G. Conditioned placebo responses. *J Pers Soc Psychol*. 1985;48(1):47-53. doi:10.1037/0022-3514.48.1.47
42. Atlas LY, Wager TD. A meta-analysis of brain mechanisms of placebo analgesia: consistent findings and unanswered questions. *Handb Exp Pharmacol*. 2014;225:37-69. doi:10.1007/978-3-662-44519-8\_3
43. Wager TD, Rilling JK, Smith EE, et al. Placebo-induced changes in fMRI in the anticipation and experience of pain. *Science*. 2004;303(5661):1162-1167. doi:10.1126/science.1093065
44. Schmid J, Theysohn N, Gaß F, et al. Neural mechanisms mediating positive and negative treatment expectations in visceral pain: a functional magnetic resonance imaging study on placebo and nocebo effects in healthy volunteers. *Pain*. 2013;154(11):2372-2380. doi:10.1016/j.pain.2013.07.013
45. Theysohn N, Schmid J, Icenhour A, et al. Are there sex differences in placebo analgesia during visceral pain processing? a fMRI study in healthy subjects. *Neurogastroenterol Motil*. 2014;26(12):1743-1753. doi:10.1111/nmo.12454
46. Kong J, Gollub RL, Rosman IS, et al. Brain activity associated with expectancy-enhanced placebo analgesia as measured by functional magnetic resonance imaging. *J Neurosci*. 2006;26(2):381-388. doi:10.1523/JNEUROSCI.3556-05.2006
47. Zeidan F, Emerson NM, Farris SR, et al. Mindfulness meditation-based pain relief employs different neural mechanisms than placebo and sham mindfulness meditation-induced analgesia. *J Neurosci*. 2015;35(46):15307-15325. doi:10.1523/JNEUROSCI.2542-15.2015
48. Kong J, Kaptchuk TJ, Polich G, et al. Expectancy and treatment interactions: a dissociation between acupuncture analgesia and expectancy evoked placebo analgesia. *Neuroimage*. 2009;45(3):940-949. doi:10.1016/j.neuroimage.2008.12.025
49. Bingel U, Wanigasekera V, Wiech K, et al. The effect of treatment expectation on drug efficacy: imaging the analgesic benefit of the opioid remifentanyl. *Sci Transl Med*. 2011;3(70):70ra14. doi:10.1126/scitranslmed.3001244
50. Kessner S, Wiech K, Forkmann K, Ploner M, Bingel U. The effect of treatment history on therapeutic outcome: an experimental approach. *JAMA Intern Med*. 2013;173(15):1468-1469. doi:10.1001/jamainternmed.2013.6705
51. Ellingsen D-M, Wessberg J, Eikemo M, et al. Placebo improves pleasure and pain through opposite modulation of sensory processing. *Proc Natl Acad Sci U S A*. 2013;110(44):17993-17998. doi:10.1073/pnas.1305050110
52. Atlas LY, Whittington RA, Lindquist MA, Wielgosz J, Sonty N, Wager TD. Dissociable influences of opiates and expectations on pain. *J Neurosci*. 2012;32(23):8053-8064. doi:10.1523/JNEUROSCI.0383-12.2012
53. Freeman S, Yu R, Egorova N, et al. Distinct neural representations of placebo and nocebo effects. *Neuroimage*. 2015;112:197-207. doi:10.1016/j.neuroimage.2015.03.015

54. Rütgen M, Seidel E-M, Silani G, et al. Placebo analgesia and its opioidergic regulation suggest that empathy for pain is grounded in self pain. *Proc Natl Acad Sci U S A*. 2015;112(41):E5638-E5646. doi:10.1073/pnas.1511269112
55. Bingel U, Lorenz J, Schoell E, Weiller C, Büchel C. Mechanisms of placebo analgesia: rACC recruitment of a subcortical antinociceptive network. *Pain*. 2006;120(1-2):8-15. doi:10.1016/j.pain.2005.08.027
56. Choi JC, Yi D-J, Han BS, Lee PH, Kim JH, Kim B-H. Placebo effects on analgesia related to testosterone and premotor activation. *Neuroreport*. 2011;22(9):419-423. doi:10.1097/WNR.0b013e32834601c9
57. Geuter S, Eippert F, Hindi Attar C, Büchel C. Cortical and subcortical responses to high and low effective placebo treatments. *Neuroimage*. 2013;67:227-236. doi:10.1016/j.neuroimage.2012.11.029
58. Wrobel N, Wiech K, Forkmann K, Ritter C, Bingel U. Haloperidol blocks dorsal striatum activity but not analgesia in a placebo paradigm. *Cortex*. 2014;57:60-73. doi:10.1016/j.cortex.2014.02.023
59. Deeks JJ, Higgins JP. Statistical algorithms in review manager 5 on behalf of the Statistical Methods Group of The Cochrane Collaboration. <http://training.cochrane.org/handbook/statistical-methods-revman5>. Published 2010. Accessed June 15, 2018.
60. Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. Effect sizes based on means. In: Borenstein M, Hedges LV, Higgins JPT, Rothstein HR, eds. *Introduction to Meta-Analysis*. Hoboken, NJ: John Wiley & Sons, Ltd; 2009. doi:10.1002/9780470743386.ch4
61. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Front Psychol*. 2013;4 (November):863. doi:10.3389/fpsyg.2013.00863
62. Higgins J, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0*. London, England: The Cochrane Collaboration; 2011.
63. Dienes Z. Using Bayes to get the most out of non-significant results. *Front Psychol*. 2014;5(July):781. doi:10.3389/fpsyg.2014.00781
64. Elsenbruch S, Kotsis V, Benson S, et al. Neural mechanisms mediating the effects of expectation in visceral placebo analgesia: an fMRI study in healthy placebo responders and nonresponders. *Pain*. 2012;153(2):382-390. doi:10.1016/j.pain.2011.10.036
65. Lui F, Colloca L, Duzzi D, Anchisi D, Benedetti F, Porro CA. Neural bases of conditioned placebo analgesia. *Pain*. 2010;151(3):816-824. doi:10.1016/j.pain.2010.09.021
66. Cohen J. A power primer. *Psychol Bull*. 1992;112(1):155-159. doi:10.1037/0033-2909.112.1.155
67. Zunhammer M, Geis S, Busch V, Greenlee MW, Eichhammer P. Effects of intranasal oxytocin on thermal pain in healthy men: a randomized functional magnetic resonance imaging study. *Psychosom Med*. 2015;77(2):156-166. doi:10.1097/PSY.0000000000000142
68. López-Solà M, Woo C-W, Pujol J, et al. Towards a neurophysiological signature for fibromyalgia. *Pain*. 2017;158(1):34-47. doi:10.1097/j.pain.0000000000000707
69. Krishnan A, Woo CW, Chang LJ, et al. Somatic and vicarious pain are represented by dissociable multivariate brain patterns. *Elife*. 2016;5:e15166. doi:10.7554/eLife.15166
70. Eippert F, Finsterbusch J, Bingel U, Büchel C. Direct evidence for spinal cord involvement in placebo analgesia. *Science*. 2009;326(5951):404. doi:10.1126/science.1180142
71. Roy M, Shohamy D, Daw N, Jepma M, Wimmer GE, Wager TD. Representation of aversive prediction errors in the human periaqueductal gray. *Nat Neurosci*. 2014;17(11):1607-1612. doi:10.1038/nn.3832
72. Schweinhardt P, Seminowicz DA, Jaeger E, Duncan GH, Bushnell MC. The anatomy of the mesolimbic reward system: a link between personality and the placebo analgesic response. *J Neurosci*. 2009;29(15):4882-4887. doi:10.1523/JNEUROSCI.5634-08.2009
73. Wechsler ME, Kelley JM, Boyd IOE, et al. Active albuterol or placebo, sham acupuncture, or no intervention in asthma. *N Engl J Med*. 2011;365(2):119-126. doi:10.1056/NEJMoa1103319
74. Benedetti F. Placebo effects: from the neurobiological paradigm to translational implications. *Neuron*. 2014;84(3):623-637. doi:10.1016/j.neuron.2014.10.023
75. Anderson SR, Losin EAR. A sociocultural neuroscience approach to pain. *Cult Brain*. 2017;5(1):14-35. doi:10.1007/s40167-016-0037-4
76. Borsook D, Becerra L, Hargreaves R. Biomarkers for chronic pain and analgesia: part 2: how, where, and what to look for using functional imaging. *Discov Med*. 2011;11(58):209-219.
77. Gatchel RJ, Peng YB, Peters ML, Fuchs PN, Turk DC. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychol Bull*. 2007;133(4):581-624. doi:10.1037/0033-2909.133.4.581
78. Ploner M, Bingel U, Wiech K. Towards a taxonomy of pain modulations. *Trends Cogn Sci*. 2015;19(4):180-182. doi:10.1016/j.tics.2015.02.007
79. Schenk LA, Sprenger C, Geuter S, Büchel C. Expectation requires treatment to boost pain relief: an fMRI study. *Pain*. 2014;155(1):150-157. doi:10.1016/j.pain.2013.09.024